



Users Online: 666

About us | Subscribe | e-Alerts | Feedback | Login

Current Issue | Archives | Ahead Of Print



Journal of Minimal Access Surgery

Search

Back

UNUSUAL CASE

Ahead of Print

Adenocarcinoma of oesophagus involving gastro-oesophageal junction following mini-gastric bypass/one anastomosis gastric bypass

Sandeep Aggarwal¹, Amit Bhambri¹, Vitish Singla¹, Nihar Ranjan Dash², Atul Sharma³

- ¹ Department of Surgical Disciplines, All India Institute of Medical Sciences, New Delhi, India
- ² Department of Gastro-intestinal Surgery, All India Institute of Medical Sciences, New Delhi, India
- ³ Department of Medical Oncology, All India Institute of Medical Sciences, New Delhi, India

Correspondence Address:

Sandeep Aggarwal,

Room No. 5034, Department of Surgical Disciplines, All India Institute of Medical Sciences, New Delhi

India

Login to access the email ID

Source of Support: None, Conflict of Interest: None

DOI: 10.4103/jmas.JMAS_320_18

PMID: 30777997

Mini-gastric bypass/one anastomosis gastric bypass (MGB/OAGB) is an emerging weight loss surgical procedure. There are serious concerns not only regarding the symptomatic biliary reflux into the stomach and the oesophagus but also the increased risk of malignancy after MGB/OAGB. A 54-year-old male, with a body mass index (BMI) of 46.1 kg/m², underwent Robotic MGB at another centre on 22nd June 2016. His pre-operative upper gastrointestinal endoscopy was not done. He lost 58 kg within 18 months after the surgery and attained a BMI of 25.1 kg/m². However, 2-year post-MGB, the patient had rapid weight loss of 19 kg with a decrease in BMI to 18.3 kg/m² within a span of 2 months. He also developed progressive dysphagia and had recurrent episodes of non-bilious vomiting. His endoscopy showed eccentric ulcerated growth in lower oesophagus extending up to the gastro-oesophageal junction and biopsy reported adenocarcinoma of oesophagus. MGB/OAGB has a potential for bile reflux with increased chances of malignancy. Surveillance by endoscopy at regular intervals for all patients who have undergone MGB/OAGB might help in early detection of Barrett's oesophagus or carcinoma of oesophagus or stomach.



Search Pubmed for

- Aggarwal S
- Bhambri A
- Singla V
- Dash NR
- Sharma A

Article Access Statistics

Reader Comments

Citation Manager

* Requires registration (Free)

Article Access Statistics

Viewed 254

PDF Downloaded 10

Recommend this journal for your library

Register with us

- > Get PDF of articles
- > Get e-mail alerts
- > Recieve notification of events
- > Add comments on articles
- > Add meetings/events

Click on image for details.

Contact us | Sitemap | Advertise with us | What's New | Disclaimer

© 2004 Journal of Minimal Access Surgery Published by Wolters Kluwer - Medknow

Online since 15th August '04

Editorial and Ethics Policies

(cc) BY-NO-SA Open Access View mobile site
ISSN: Print -0972-9941, Online - 1998-3921

