

[J Minim Access Surg](#). 2017 Oct-Dec; 13(4): 291–295.

PMCID: PMC5607797

doi: [10.4103/jmas.JMAS_5_17](https://doi.org/10.4103/jmas.JMAS_5_17)

PMID: [28872099](https://pubmed.ncbi.nlm.nih.gov/28872099/)

Impact of bariatric surgery on obstructive sleep apnoea–hypopnea syndrome in morbidly obese patients

[Pratyusha Priyadarshini](#), [Vijay Pal Singh](#), [Sandeep Aggarwal](#), [Harshit Garg](#), [Sanjeev Sinha](#),¹ and [Randeep Guleria](#)²

Department of Surgical Disciplines, All India Institute of Medical Sciences, New Delhi, India

¹Department of Medicine, All India Institute of Medical Sciences, New Delhi, India

²Department of Pulmonary Medicine and Sleep Disorders, All India Institute of Medical Sciences, New Delhi, India

Address for correspondence: Dr. Sandeep Aggarwal, Department of Surgical Disciplines, All India Institute of Medical Sciences, Ansari Nagar, New Delhi, India. E-mail: sandeep_aiims@yahoo.co.in

Received 2017 Jan 11; Accepted 2017 Mar 2.

[Copyright](#) : © 2017 Journal of Minimal Access Surgery

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Abstract

Background:

Obstructive sleep apnea (OSA) is commonly associated with morbid obesity. Weight loss following bariatric surgery results in resolution or improvement of OSA. However, few studies have done objective assessment of the impact of bariatric surgery on OSA.

Objective:

The aim of this study was to assess the outcome of bariatric surgery on OSA.

Setting:

The study was conducted in the teaching institution of a tertiary care centre.

Methods:

Twenty-seven morbidly obese patients seeking bariatric surgery were administered Epworth Sleepiness Scale (ESS) health questionnaire and subjected to overnight polysomnography. Repeat assessment using ESS and polysomnography was done at 3–6 months after surgery.

Results:

Mean age was 42.4 ± 10.5 years, and majority (77.8%) were female. The mean pre-operative weight and body mass index (BMI) were 126.4 ± 24.9 kg and 48.4 ± 8.2 kg/m², respectively. Nearly 29.6% patients had symptoms of excessive daytime somnolence based on ESS score and overnight polysomnography detected the presence of OSA in 96.3% patients, of which 51.9% had severe OSA. At mean follow-up of 5.2 ± 2.5 months after surgery, mean weight and BMI decreased to 107.4 ± 24.5 kg and 41.2 ± 8.2 kg/m², respectively. Mean ESS score and mean apnoea–hypopnea index declined